



# In vivo fiber tractography of the right and left ventricles using diffusion tensor MRI of the entire human heart

## Citation

Mekkaoui, Choukri, Timothy G Reese, Marcel P Jackowski, Himanshu Bhat, William J Kostis, and David E Sosnovik. 2014. "In vivo fiber tractography of the right and left ventricles using diffusion tensor MRI of the entire human heart." *Journal of Cardiovascular Magnetic Resonance* 16 (Suppl 1): P17. doi:10.1186/1532-429X-16-S1-P17. <http://dx.doi.org/10.1186/1532-429X-16-S1-P17>.

## Published Version

doi:10.1186/1532-429X-16-S1-P17

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12406803>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



**POSTER PRESENTATION**

**Open Access**

# *In vivo* fiber tractography of the right and left ventricles using diffusion tensor MRI of the entire human heart

Choukri Mekkaoui<sup>1\*</sup>, Timothy G Reese<sup>1</sup>, Marcel P Jackowski<sup>2</sup>, Himanshu Bhat<sup>3</sup>, William J Kostis<sup>1</sup>, David E Sosnovik<sup>1</sup>

From 17th Annual SCMR Scientific Sessions  
New Orleans, LA, USA. 16-19 January 2014

## Background

Diffusion Tensor MRI (DTI) tractography of the human heart *in vivo* has previously been performed with either large slice gaps or limited coverage [1,2]. The aim of this study was to investigate the feasibility of performing DTI of the entire human heart *in vivo* without slice gaps. This, we hypothesized, would enhance the characterization of fiber architecture in the left ventricle (LV), allow myofiber organization in the right ventricle (RV) to be characterized *in vivo*, and further elucidate microstructural differences in the heart between systole and diastole.

## Methods

DTI was performed on a clinical 3T scanner (Skyra, Siemens) using a fat-suppressed, zone-selected, diffusion-encoded stimulated echo sequence with 10 diffusion encoding directions, TE/TR 33/80 ms, GRAPPA rate 2, b-value 500 s/mm<sup>2</sup>, resolution 2.8 × 2.8 × 8 mm<sup>3</sup>, 8 averages and multiple breath-holds. The entire LV and RV were covered in 13 contiguous short-axis slices. Images were acquired in the systolic and diastolic sweet spots [3] of the cardiac cycle and were spatiotemporally coregistered [4]. Tractography was performed by numerically integrating the primary eigenvector field into streamlines using an adaptive 5<sup>th</sup> order Runge-Kutta method. The impact of cumulative image averages (1-8) on the reliability of the fractional anisotropy (FA) and fiber helix angle (HA) indices was assessed.

## Results

A composite view of the anterior thorax and heart is shown in Figure 1A. The contrast between the helical

pattern of the fibers in the heart (color-coded by HA) and the linear organization of skeletal muscle fibers in the chest wall can be seen. The RV consisted of a bilayer of obliquely-oriented fibers, lacking circumferential fibers (Figure 1B). In contrast, fibers in the midwall of the LV were circumferential, forming a distinct band between the subendocardial and subepicardial fibers (Figure 1C). Figure 1D shows the tractogram of the entire heart, depicting the intertwined arrangement of myofibers at the anteroseptal RV-LV junction. Figure 2A and 2B show that FA values in both sweet spots were significantly overestimated when the number of averages used was < 5 ( $p < 0.01$ ). In contrast, convergence to a stable FA value (LV, diastole:  $0.42 \pm 0.03$ , systole:  $0.41 \pm 0.03$ ; RV, diastole:  $0.32 \pm 0.11$ , systole:  $0.36 \pm 0.02$ ) was observed from 5-8 averages. The transmural gradient in HA exhibited a similar pattern of convergence but required more averages (Figure 2C, D).


## Conclusions

DTI-tractography of the entire human heart can be performed *in vivo*, without slice gaps, and allows both the LV and the RV to be characterized. A minimum of 5 averages at each slice location is required for accurate quantification. The ability to characterize fiber architecture in the LV and RV *in vivo* has the potential to provide new insights into a range of diseases affecting both the pulmonary venous and arterial circulations.


## Funding

R01 HL093038 (D.E.S.), R01HL112831 (D.E.S.), P41RR14075 (Martinos Center).

<sup>1</sup>Radiology, Harvard Medical School - Massachusetts General Hospital -  
Martinos Center for Biomedical Imaging, Charlestown, Massachusetts, USA  
Full list of author information is available at the end of the article



**Figure 1** (A) A composite view of the anterior thorax and heart shows the contrast between the helical pattern of the fibers in the myocardium (color-coded by HA) and the linear organization of skeletal muscle fibers in the chest wall. (B) The fiber architecture of the RV can be well resolved *in vivo* and lacks the circumferential fibers seen in the LV (C). (D) Tractography within a slab located at the basal-level depicting the transmural myofiber arrangement of both RV and LV. The intertwined arrangement of myofibers at the antero-septal RV-LV junction is slightly magnified.



**Figure 2** (A, B) Cumulative averages (1-8) of FA for the LV and the RV, respectively. (A) In the LV, FA values stabilize after 5 averages in diastole as well as in systole. (B) In the RV, FA also stabilizes after 5 averages, albeit with a higher standard deviation in diastole, but FA values are 9% lower in diastole than in systole. (C, D) Transmural distribution of HA for 1, 4, and 8 cumulative averages in diastole and systole, respectively. For optimal estimation of HA, 8 averages are required since the HA is a function of the primary eigenvector, and more degrees of freedom need to be estimated.

#### Authors' details

<sup>1</sup>Radiology, Harvard Medical School - Massachussets General Hospital - Martinos Center for Biomedical Imaging, Charlestown, Massachusetts, USA.  
<sup>2</sup>Department of Computer Science, University of São Paulo, São Paulo, São Paulo, Brazil. <sup>3</sup>Siemens Medical Solutions USA Inc., Charlestown, Massachusetts, USA.

Published: 16 January 2014

#### References

1. Toussaint, et al: *MEDIA* 2013.
2. Nielles-Vallés, et al: *MRM* 2012.
3. Tseng, et al: *MRM* 1999.
4. Mekkaoui, et al: *JCMR* 2012.

doi:10.1186/1532-429X-16-S1-P17

**Cite this article as:** Mekkaoui et al.: *In vivo fiber tractography of the right and left ventricles using diffusion tensor MRI of the entire human heart.* *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P17.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

